

2022-11-21

# Prospects for soilless farming in Africa: A review on the aids of plant growth-promoting rhizobacteria inoculants in hydroponics

Margaret, Gumisiriza

International Journal of Biosciences

---

<http://dx.doi.org/10.12692/ijb/21.5.273-282>

*Provided with love from The Nelson Mandela African Institution of Science and Technology*



## REVIEW PAPER

## OPEN ACCESS

## Prospects for soilless farming in Africa: A review on the aids of plant growth-promoting rhizobacteria inoculants in hydroponics

Gumisiriza S. Margaret<sup>\*1</sup>, Aloo N. Becky<sup>2</sup>, Makumba Billy<sup>3</sup>, Tumuhairwe J. Baptist<sup>4</sup>, Ndakidemi A. Patrick<sup>1</sup>, Mbega R. Ernest<sup>1</sup>

<sup>1</sup>*School of Life Sciences and Bioengineering, The Nelson Mandela African Institution of Science and Technology, Arusha, Tanzania*

<sup>2</sup>*School of Science, University of Eldoret, Eldoret, Kenya*

<sup>3</sup>*School of Science, Moi University, Eldoret, Kenya*

<sup>4</sup>*Makerere University, College of Agricultural and Environmental Sciences, Department of Agricultural Production, Kampala, Uganda*

**Key words:** Hydroponics, Nutrient solubilization, Soilless culture, PGPR, Urban farming

<http://dx.doi.org/10.12692/ijb/21.5.273-282>

Article published on November 21, 2022

### Abstract

Due to the growing population in Africa, there is need to identify farming systems that can increase food productivity for the increasing population. Hydroponic farming presents a viable option for sustainable and climate resilient agricultural production especially in areas faced with environmental challenges such as; limited arable land and has the ability to realize global food security the rising population and urbanization in Africa. However, the supply of plant nutrients in sufficient and sustainable quantities at affordable costs is one of the critical and limiting factors in adoption of hydroponics. Chemical hydroponic fertilizers that are often used are not only environmentally unfriendly but also costly and less-readily available. As such, alternative crop fertilization mechanisms like the use of plant growth-promoting rhizobacteria (PGPR) as inoculants in hydroponics are not only an environmentally feasible but also economical solution for countries in Africa. There are numerous studies regarding this crop-fertilization mechanism, but these mainly refer to controlled environment such as; green-houses, and field tests. Their use in hydroponic farming is still largely unexploited. This review highlights the nutrient requirements, types and benefits of hydroponics in addition to unraveling the potential that PGPR inoculants hold as a sustainable and economical fertilizer alternative in hydroponics.

\* **Corresponding Author:** Gumisiriza S. Margaret ✉ [m5gundas@gmail.com](mailto:m5gundas@gmail.com)

## Introduction

There has been increasing rates of population across the globe of which majority of these people move to urban centres due to various reasons such as: search for employment opportunities. This increasing population around urban areas which has resulted into the growing food demand especially in under-developed states has roused interest in alternative crop farming systems due to the reducing arable land vis-a-vis the increasing food demands.

It is worth noting that crop cultivation has over time heavily relied on conventional agricultural practices which unfortunately has stimulated the emission of major greenhouse gases (GHGs) like Nitrous oxide (Di Benedetto *et al.*, 2017). Accompanying this, is the deterioration in arable land because of industrialization, urbanization and increase in population (Hlophe *et al.*, 2019; Mgbemene *et al.*, 2016) especially in African urban settings. Furthermore, soil-based farming is constantly affected by pests and diseases, and the effects of climate change, for example, water scarcity and flooding resulting from low and heavy rainfalls (Depardieu *et al.*, 2016).

Researchers presume that the solution to these challenges could be realized with the adoption of environmentally friendly crop cultivation techniques such as; hydroponic farming (Aini *et al.*, 2019). This soil-less farming system does not only improve the efficiency of land, water, and nutrients usage but also the quality and quantity of plant yields (Barbosa *et al.*, 2015).

One critical factor in hydroponic farming is the effective supply and management of plant nutrients (Aini *et al.*, 2019). Chemical fertilizers that are normally used in hydroponic farming are often costly, unsustainable and contribute to the emission of GHGs (Amgai *et al.*, 2017; Di Benedetto *et al.*, 2017) thus calling for alternative options. One of such options are; the PGPR that have also been associated with improved plant growth, facilitation of plant nutrient availability and uptake in addition to synthesis of plant growth-promoting (PGP) hormones in the rhizosphere (Ahemad and Kibret, 2014; Hayat

*et al.*, 2010). These microorganisms have been used in agriculture as bio fertilizers, bio-control agents, and bioremediation both in soil and soil-less (hydroponic) systems (Lee and Lee, 2015). However, most of these studies have dwelled on conventional farming in soils, while little is known about their use in hydroponic systems (Paradiso *et al.*, 2017). The aim of this review is to unravel the potential that PGPR have as inoculants in hydroponic farming.

The review revisits hydroponics as a climate-smart agricultural option, pinpoints some of the shortcomings and evaluates the potential of PGPR inoculants in the provision of essential nutrients in these systems while promoting sustainable agriculture. Some of the advances that have been made concerning rhizobacterial inoculation in hydroponic systems are also articulated. Finally, the review assesses the prospects of these inoculants in hydroponic systems and the overall contribution of their usage in soil-less farming as an option for sustainable agriculture and environmental management.

## Materials and methods

Literature search and assessment was done using online data sources including: Web of science, Science Direct, Scopus data base and Google scholar. The key words used for the review of articles upto 2022 data included; vegetable farming, hydroponics, soilless farming, PGPR inoculants, hydroponics and sustainable agriculture

## Discussion

### *What is hydroponics*

Hydroponic farming is a method of growing crops under soil-less conditions by dipping the roots in a water solution composed of chemical nutrients to support plant growth (Nisha *et al.*, 2018; (Barbosa *et al.*, 2015). The technology highly favors crop production in areas with non-arable soils and offers more efficient use of resources like water which undoubtedly highlights the advantages of hydroponic farming (Kinoshita *et al.*, 2016). The efficiency and effectiveness in supply of plants nutrients in hydroponics is critical to plant growth. Nitrogen (N), Phosphorus (P), Potassium (K), Calcium (Ca) and

Magnesium (Mg) are considered as the primary nutrients required for plant growth (Maneejantra *et al.*, 2016) while Manganese (Mn), Iron (Fe), Zinc (Zn), Boron (B), and Copper (Cu) are the supplementary micronutrients (Fig.1) which support the healthy growth of plants under hydroponic farming. The composition of these nutrients plays a crucial role in determining the electrical conductivity (EC) of the solution.


The Potential of Hydrogen (pH) of the nutrient solutions is also a vital factor in hydroponic farming where the solution must contain ions that can be absorbed by the plants. A pH range of 5.5-6.5 is generally ideal for nutrient availability in these farming systems but this can keep fluctuating as the crop grows and different crops have different pH requirements

(Alexopoulos *et al.*, 2021). With this agricultural technology, crops can be grown with or without the support of a medium (Fig.1) that not only acts as a conduit for nutrients and water but also offers plant support (Gumisiriza *et al.*, 2020). Fig. 1 summarizes the nutrients and media considered beneficial for effective hydroponic farming while Table 1 describes the different farming systems under hydroponics.

Primary nutrients	Secondary nutrients	Non-organic medium	Organic medium
<input type="checkbox"/> Nitrogen	<input type="checkbox"/> Iron	<input type="checkbox"/> Rock wool	<input type="checkbox"/> Sugar cane bagasse
<input type="checkbox"/> Phosphorus	<input type="checkbox"/> Manganese	<input type="checkbox"/> Vermiculite	<input type="checkbox"/> Rice hulls
<input type="checkbox"/> Potassium	<input type="checkbox"/> Boron	<input type="checkbox"/> Perlite	<input type="checkbox"/> Coco-coir
<input type="checkbox"/> Calcium	<input type="checkbox"/> Zinc	<input type="checkbox"/> Gravel	<input type="checkbox"/> Coffee husks
<input type="checkbox"/> Magnesium	<input type="checkbox"/> Copper	<input type="checkbox"/> Peat moss	<input type="checkbox"/>
		<input type="checkbox"/> Clay granules	<input type="checkbox"/>

**Fig. 1.** Summary of nutrients and essential medium for hydroponic farming.

**Table 1.** Forms of hydroponic farming.

	System	Mode of flow	Description	Benefits	Drawbacks	References
	Wick system (Passive technique)	Non-circulating system	This system uses a wick to draw the required nutrients from the nutrient reservoir into the growing medium.	Simple to build. It is cheap.	Favors small plants with low nutrient necessities. No recirculation of nutrients. Susceptible to algae.	(Nisha, 2018)
	Deep Water Culture (Direct Water Culture)	Non-circulating system	The plants are put in small baskets (net cups) fitted with growing medium and roots are suspended directly in a highly oxygenated nutrient solution.	Easy to build and operate. Presence of enough dissolved energy. It is cheap.	Roots are prone to rotting if not cleaned often. Requires solution refilling. Slow rate of growth.	(Nisha, 2018)
	Ebb and Flow (Flood and Drain)	Circulating system	The system works by flooding the plant plate with a nutrient solution using a pump that is connected to the solution tank. The solution is pumped at given time intervals with the use of a timer and then drained back to the nutrient tank.	Low maintenance costs.	Susceptible to algae. Malfunctions can lead to crop failure. Prone to blockage.	(Seungjun Lee and Jiyoung Lee, 2015)
	Nutrient film technique (NFT)	Circulating system	Plant roots are supplied with nutrients by dipping them in channels filled with the nutrient-rich solution and the system doesn't need a timer.	Plant roots receive enough nutrients, water, and oxygen. No need for a timer.	Malfunctions can lead to crop failure. Prone to blockage.	(Domingues <i>et al.</i> , 2012; Jones, 2016; Mamta and Shraddha, 2013; Omics, 2017; Wilcox, 1982)
High technology & cost	Aeroponics	Circulating system	Plant roots are suspended in air and nutrients are supplied to them in form of mist with the use of a timed pump which ensures release of mist after every few minutes.	Does not require any growing medium. Ensures adequate nutrient absorption.	Pump interference can lead to root drying. High and expensive technology. Consumes time. Malfunctions can lead to crop failure.	(Mazhar, 2020)

#### Benefits of hydroponic farming

Hydroponic farming has a number of benefits over conventional farming. It is more efficient in water

utilization and favors the production of high and consistent yields in soil-less areas (Gruda., 2019; Gumisiriza *et al.*, 2022).

This farming system is also less labor-intensive as it requires no weeding and land preparation like most conventional farming practices (Pignata *et al.*, 2017). According to several researchers, hydroponic systems may also be used to improve vegetables and fruits both in terms of nutrition, quality, and shelf life according to the market and consumer needs (Amalfitano *et al.*, 2017; Buchanan and Omaye, 2013; Islam *et al.*, 2018; Selma *et al.*, 2012; Sgherri *et al.*, 2010). There is reduced use of pesticides and fungicides since the system is often practiced under a climate-controlled environment (Benke and Tomkins, 2017). Despite the mentioned benefits, the technology requires adequate technical knowledge and high investment costs which include; PVC pipes, timers, hydroponic fertilizers etc... (Nisha, 2018; Gumisiriza *et al.*, 2022a). Hydroponic systems can also be prone to pathogens, and fungal infections (Constantino *et al.*, 2013; Li *et al.*, 2014; Song *et al.*, 2004). The media used in hydroponic farming to deliver the nutrients to the plant roots are not rich with microorganisms that can boost the availability and uptake of nutrients by plants (Hatice *et al.*, 2012).

#### *Rhizobacteria and plant growth promotion*

Rhizobacteria are plant root-colonizing bacteria that can stimulate plant development by promoting root growth (Grover *et al.*, 2011). For several decades, researchers have highlighted the importance of PGPR in plant mineral nutrition (Pii *et al.*, 2015). The Food and Agriculture Organisation (FAO) estimates that the demand for Nitrogen fertilizers will exceed 130 million tons per year which is environmentally unsuitable especially since their production largely depends on the use of fossil fuels (Kliopova *et al.*, 2016). Many PGPR have attracted the attention of researchers as plant inoculants due to their capacity to increase nutrient uptake in crops by the production of PGP hormones like indole acetic acid (IAA) and Gibberellic acids (GA) (Choudhary *et al.*, 2018; Vejan *et al.*, 2016). These hormones have been associated with increased foliage and root elongation (Hassen *et al.*, 2016; Vacheron *et al.*, 2015).

There is no doubt that these bacterial substances can significantly increase crop yields and open up a new

horizon for sustainable plant productivity. Iron (Fe) is an important micronutrient needed for plant growth (Saha *et al.*, 2016). It's unavailability is a major plant-growth limiting factor in crop production systems (Arora and Verma, 2017; Singh *et al.*, 2019).

Some bacteria have special mechanisms for Fe-acquisition by synthesizing low molecular weight metabolites known as siderophores (Maheshwari *et al.*, 2019) with high affinity for Fe in low-Fe conditions (Mhlongo *et al.*, 2018; Tank *et al.*, 2012). This way, the siderophores function as Fe-chelators and bind most of the available Fe in the rhizosphere (Singh *et al.*, 2019). Furthermore, literature advances that siderophore-producing bacteria and the subsequent Fe-unavailability in plant rhizospheres may also prevent the proliferation of plant pathogens (Mitter *et al.*, 2013; Olanrewaju *et al.*, 2017).

A lot of studies have shown the ability of different rhizobacterial species to produce siderophores and the enhancement of Fe nutrition in different crops (Emami *et al.*, 2019; Ghavami *et al.*, 2017; Liaqat and Eltem, 2016). Siderophore production is a typical example of how rhizobacterial inoculants can establish themselves in the rhizosphere and enhance Fe nutrition. Due to its indisputable importance, it should be given more attention (Aloo *et al.*, 2019).

#### *Advances in hydroponic vegetable production using rhizobacterial inoculation*

Research studies have shown instances where the use of PGPR inoculants has demonstrated positive effects on plant growth in hydroponics. For instance, lettuce seedlings grew well when microbial culture solution was used in hydroponics (Shinohara *et al.*, 2011).

Some PGPR like *Pseudomonas* spp., *Bacillus* spp. have successfully improved the growth, yield, and quality of vegetables like; cucumber, lettuce, and tomato under hydroponic farming (Lee and Lee, 2015). In a study by Pii *et al.*, 2018).

Other instances where rhizobacterial inoculations have successfully been shown to promote the growth of different vegetables under hydroponic farming are summarized in Table 2.

**Table 2.** Studies demonstrating the growth promotion of different vegetables using rhizobacterial inoculants in hydroponic systems.

Crop	Inoculant	Effects on growth	Reference
Tomato and pepper	<i>Bacillus licheniformis</i>	Increased fruit yield	(Garcia <i>et al.</i> , 2004)
Tomato	<i>Pseudomonad fluorescens</i> , <i>Serratia marcescens</i> , <i>Bacillus</i> sp.	Increased fruit yield Increased water efficiency	(Kıdoğlu <i>et al.</i> , 2009)
Tomato	<i>P. fluorescens</i> , <i>P. Putida</i>	Increased weight and number of fruits	(Gul <i>et al.</i> , 2012)
Lettuce	<i>Pseudomonas</i> sp.	Increased root and shoot mass	(Cipriano <i>et al.</i> , 2014; Tahmasbi <i>et al.</i> , 2014)
Tomato, cucumber, lettuce, and Irish potato	<i>Pseudomonas</i> sp.	Increased shoot and root weight	(Peer and Schippers, 1989)
Tomato	<i>P. fluorescens</i> , <i>Burkholderia cepacian</i> , <i>Stenotrophomonas maltophilia</i> , ( <i>P. fluorescens</i> , <i>S. maltophilia</i> )	Rapid growth and increased number of leaves	(Alsanius and Gertsson, 2004)
Tomato	<i>Azotobacter chroococcum</i> , <i>Azospirillum brasilense</i> , <i>Pseudomonas flourescens</i> , and <i>Bacillus subtilis</i>	Increased yield and quality	(Aini <i>et al.</i> , 2019)
Tomato	<i>Not mentioned</i>	Increased growth biometrics	(Dasgan <i>et al.</i> , 2017)
Lentils	<i>Rhizobium</i> strains		(Zafar <i>et al.</i> , 2012)

While testing various commercially-available bio fertilizers containing various rhizobacterial inoculants like; *B. subtilis*, *Azotobacter vinelandi* and *Clostridium pasteurianum* to replace chemical fertilizers in hydroponically grown squash (*Cucurbita pepo*) in Turkey, (Dasgan *et al.*, 2012), significant increases in leaf area and nutrient uptake were observed in the bio fertilizer-treated plants than the chemically-treated plants.

It was concluded that the bio-fertilizers could be used in soilless squash production to reduce chemical fertilization resulting in less environmental impact. For years, several efforts have been made to demonstrate endophytic and associative N<sub>2</sub> fixation in crops using free-living diazotrophs (Ahemad and Kibret, 2014; da Silva *et al.*, 2012; Santoyo *et al.*, 2016). However, the contribution of symbiotically-fixed N to hydroponically-grown plants remains largely unestablished and deficient. Previous studies on hydroponically-grown soybean have demonstrated BNF by *Ochrobactrum* bacteria (Paradiso *et al.*, 2017; Paradiso *et al.*, 2015; Paradiso *et al.*, 2014), where inoculation seemed to promote the growth of root system, likely enhancing the capability of nutrient uptake. The use of rhizobacterial inoculants in soilless

culture is becoming more and more common (Paradiso *et al.*, 2017).

In a study by Phibunwatthanawong and Riddech (2019) on the formulation of liquid organic fertilizers for growing vegetables under hydroponic conditions using rhizobacterial inoculants, it was shown that all formulations contained IAA whose quantities increased with incubation up to 59.53mg L<sup>-1</sup>. The production of IAA, increased grain yield, root length and number of nodules per plant has also been shown under hydroponic farming conditions (Zafar *et al.*, 2012). The production of IAA and other PGP hormones has especially been linked to the increased uptake of different nutrients by plants.

#### *The future prospects of rhizobacteria inoculation in hydroponic systems*

Rhizobacterial inoculants are advantageous and unique forms of organic fertilizers because of their abilities to synthesize PGP hormones that do not occur in chemical fertilizers (Phibunwatthanawong and Riddech, 2019). However, majority of the studies related to their importance in plant growth have been carried out in soil-based systems, while very little is known about their potential in hydroponic systems



where their efficiency also depends on the ability to proliferate and colonize plant roots (Lee and Lee, 2015). The experimental evidence on their efficacy at the field level is still grossly inadequate.

This calls for more research to reveal their efficiency and increase their usability in hydroponic systems. This and related information will certainly help in understanding their use as bio inoculants for practical purposes (Teotia *et al.*, 2016). The productivity and quality of crops grown in hydroponic systems are markedly dependent on the adequate supply, availability and uptake of plant nutrients (Valentinuzzi *et al.*, 2015). The mechanisms underlying nutrition in plants, their regulation and the bio-geochemical cycles of nutrients in hydroponic systems need to be understood to fully develop PGPR strategies for the optimization of the hydroponic crop cultivation (Sambo *et al.*, 2019).

Biological nitrogen fixation offers a viable option for minimizing the use of chemical fertilizers in hydroponic systems. As such, more research in this area is required for vegetables to promote this option for sustainable agriculture especially in low developed countries such as; Africa. This will also play a huge role in reducing the utilization of chemical fertilizers and artificial growth regulators in hydroponic systems (Prathap and Ranjitha, 2015) as well as increase food production to meet the demands of the growing population especially in urban centres.

### Concluding remarks and recommendations

Hydroponics is an evolving field of urban agriculture especially in Africa that presents an ultimate solution to many of the contemporary challenges of soil-based farming. Nevertheless, the provision of appropriate nutrients at a low cost and in an organic and environmentally-friendly manner remains a huge hinderance in the adoption of this farming system. Some studies have explored the role of organic fertilizers like animal manures such as; bat, pig, cow, fish and chicken dung in hydroponic farming. However, only few have considered the use of rhizobacteria yet these can also improve plant nutrient availability and uptake in these systems.

There is need for more research into this aspect to lower the costs of hydroponic farming in terms of hydroponic fertilizers, reduce the environmental impacts associated with chemical fertilizers and promote organic hydroponic farming. These and more initiatives can eventually increase the adoption of this technology for urban agriculture as well as environmental sustainability. This will also address the increasing concerns across the globe on whether hydroponics is an organic farming system by highlighting the role of rhizobacteria as an organic fertilizer substitute in hydroponic farming.

### Declaration of competing interest

The authors declare that they have no conflict of interest.

### References

- Ahemad M, Kibret M.** 2014. Mechanisms and applications of plant growth promoting rhizobacteria: Current perspective. *Journal of King Saud University-Science* **26**, 1-20.
- Aini N, Yamika WSD, Pahlevi RW.** 2019. The effect of nutrient concentration and inoculation of PGPR and AMF on the yield and fruit quality of hydroponic cherry tomatoes (*Lycopersicon esculentum* Mill. Var. Cerasiforme). *Journal of Applied Horticulture* **21**, 116-122.
- Aloo BN, Mbega ER, Makumba BA.** 2019. Rhizobacteria-Based Technology for Sustainable Cropping of Potato (*Solanum tuberosum* L.). *Potato Research* 1-21.
- Alsanius BW, Gertsson UE.** 2004. Plant Response of Hydroponically Grown Tomato to Bacterization. *Acta Horticulturae* **644**, 583-588.
- Amalfitano C, Del Vacchio L, Somma S, Cuciniello A, Caruso G.** 2017. Effects of cultural cycle and nutrient solution electrical conductivity on plant growth, yield and fruit quality of "Friariello" pepper grown in hydroponics. *Horticultural Science* **44**, 91-98.
- Angai S, Paudel SR, Bista DR, Poudel SR.** 2017. Government intervention on organic fertilizer promotion: A key to enhancing soil health and environment. *The Journal of Agriculture and Environment* **18**, 131-138.

- Arora NK, Verma M.** 2017. Modified microplate method for rapid and efficient estimation of siderophore produced by bacteria. *BioTechnology* **7**, 381.
- Barbosa GL.** 2015. Comparison of land, water, and energy requirements of lettuce grown using hydroponic vs. Conventional agricultural methods. *International Environmental Research and Public Health* **1**, 6879-6891.
- Benke K, Tomkins B.** 2017. Future food-production systems: vertical farming and controlled-environment agriculture. *Sustainability Science* **13**, 13-26.
- Buchanan DN, Omaye ST.** 2013. Comparative Study of Ascorbic Acid and Tocopherol Concentrations in Hydroponic- and Soil-Grown Lettuces. *Food and Nutrition Sciences* **04**, 1047-1053.
- Choudhary M.** 2018. Towards Plant-Beneficiary Rhizobacteria and Agricultural Sustainability. Role of Rhizospheric Microbes in Soil: Volume 2: Nutrient Management and Crop Improvement **2**, 1-46.
- Cipriano MAP, Patricio FRA, Freitas SS.** 2014. Potential of rhizobacteria to promote root rot growth and control in hydroponically cultivated lettuce. *Summa Phytopathologica* **39**, 51-57.
- Constantino NN.** 2013. Root-expressed maize lipoxygenase 3 negatively regulates induced systemic resistance to *Colletotrichum graminicola* in shoots. *Front Plant Science* **4**.
- da Silva MF.** 2012. Survival of endophytic bacteria in polymer-based inoculants and efficiency of their application to sugarcane. *Plant and Soil* **356**, 231-243.
- Dasgan HY, Aydoner G, Akyol M.** 2012. Use of some micro-organisms as bio-fertilizers in soilless grown squash for saving chemical nutrients. *Acta Horticulturae* **927**, 155-162.
- Dasgan HY, Cetinturk T, Altuntas O.** 2017. The effects of biofertilisers on soilless organically grown greenhouse tomato. *Acta Horticulturae* **1164**, 555-561.
- Depardieu C, Prémont V, Boily C, Caron J.** 2016. Sawdust and Bark-Based Substrates for Soilless Strawberry Production: Irrigation and Electrical Conductivity Management. *PLoS ONE* **11**, 4.
- Di Benedetto NA.** 2017. The role of Plant Growth Promoting Bacteria in improving nitrogen use efficiency for sustainable crop production: A focus on wheat. *AIMS Microbiology* **3**, 413-434.
- Domingues DS, Takahashi HW, Camara CAP, Nixdorf SL.** 2012. Automated system developed to control pH and concentration of nutrient solution evaluated in hydroponic lettuce production. *Computer Electronic Agriculture* **84**, 53-61.
- Emami S, Alikhani HA, Pourbabaei AA, Etesami H, Motessharezadeh B.** 2019. Effect of rhizospheric and endophytic bacteria with multiple plant growth promoting traits on wheat growth. *Environmental Science Pollution Research* **26**, 29804.
- Garcia JAL, Ramos APB, Palomino M, Manero FJG.** 2004. Effect of inoculation of *Bacillus licheniformis* on tomato and pepper. *Agronomie* **24**, 169-176.
- Ghavami N, Alikhani HA, Pourbabeii AA, Besharati H.** 2017. Effects of two new siderophore-producing rhizobacteria on growth and iron content of maize and canola plants. *Journal of Plant Nutrition* **40**, 736-746.
- Grover M, Ali SKZ, Sandhya V, Rasul A, Venkateswarlu B.** 2011. Role of microorganisms in adaptation of agriculture crops to abiotic stresses. *World Journal of Microbiology and Biotechnology* **27**, 1231-1240.
- Gul A.** 2012. Effect of Rhizobacteria on Yield of Hydroponically Grown Tomato Plants. *Acta Horticulturae* **952**, 777-784.
- Hassen AI, Bopape FL, Sanger LK.** 2016. Microbial inoculants as agents of growth promotion and abiotic stress tolerance in plants. *Microbial inoculants in sustainable agricultural productivity* Springer 23-36.



- Hatice O, Birsen C, Lalehan Y, Muge Sahin.** 2012. Effect of Rhizobacteria on Hydroponically grown tomato plants. *Acta Horticulture* **952**, 777-784.
- Hayat R, Ali S, Khalid R, Ahmed I.** 2010. Soil beneficial bacteria and their role in plant growth promotion: A review. *Annals of Microbiology* **60**, 579-598.
- Hlophe P.** 2019. Effects of different media on the growth and yield of Swiss chard (*Beta vulgaris* var. *cicla*) grown in hydroponics. *Horticulture International Journal* **3**, 147-151
- Islam MZ, Mele MA, Baek JP, Kang H.** 2018. Iron, iodine and selenium effects on quality, shelf life and microbial activity of cherry tomatoes. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca* **46**, 388-392.
- Jones J.** 2016. *Hydroponics: A Practical Guide for the Soilless Grower*. CRC Press., Boca Raton.
- Kıdoğlu F, Gül A, Tüzel Y, Özaktan H.** 2009. Yiled Enhancement of Hydroponically-Grown Tomatoes by Rhizobacteria. *Acta Horticulturae* **807**, 457-480.
- Kinoshita T, Yamazaki H, Inamoto K, Yamazaki H.** 2016. Analysis of yield components and dry matter production in a simplified soilless tomato culture system by using controlled-release fertilizers during summer-winter greenhouse production. *Science and Horticulture* **202**, 17-24.
- Kliopova I, Baranauskaite-Fedorova I, Malinauskiene M, Staniškis K.** 2016. Possibilities of increasing resource efficiency in nitrogen fertilizer production. *Clean Technologies and Environmental Policy* **18**, 901-914.
- Lee SW, Lee J.** 2015. Beneficial bacteria and fungi in hydroponic systems: Types and characteristics of hydroponic food production methods. *Science and Horticulture* **195**, 206-215.
- Li M.** 2014. Monitoring by real-time PCR of three water-borne zoospore Pythium species in potted flower and tomato greenhouses under hydroponic culture systems. *European Journal of Plant Pathology* **140**, 229-242.
- Liaqat F, Eltem R.** 2016. Identification and Characterization of endophytic bacteria isolated from in vitro cultures of peach and pea rootstocks. *BioTechnology* **6**, 2-9.
- Maheshwari R, Bhutani N, Suneja P.** 2019. Screening and Characterization of siderophore producing endophytic bacteria from *Cicer arietinum* and *Pisum sativum* plants *Journal of Applied Biology and Biotechnology* **7**, 7-14.
- Mamta Shraddha.** 2013. A review on plant without soil -Hydroponics *International Journal of Research in Engineering and Technology* **2**.
- Maneejantra N, Tsukagoshi S, Lu N, Supaibulwatana K, Takagaki M.** 2016. A Quantitative Analysis of Nutrient Requirements for Hydroponic Spinach (*Spinacia oleracea* L.) Production Under Artificial Light in a Plant Factory. *Journal of Fertilizers and Pesticides* **7**, 2471-2728.
- Mgbemene CA, Mnaji CC, Nwozor C.** 2016. Industrialization and its Backlash: Focus on Climate Change and its Consequences. *Journal of Environmental Science and Technology* **9**, 301-316.
- Mhlongo MI, Piater LA, Madala NE, Labuschagne N, Dubery IA.** 2018. The Chemistry of Plant-Microbe Interactions in the Rhizosphere and the Potential for Metabolomics to Reveal Signaling Related to Defense Priming and Induced Systemic Resistance. *Frontiers in Plant Science* **9**, 112.
- Mitter B.** 2013. Comparative genome analysis of *Burkholderia phytofirmans* PsJN reveals a wide spectrum of endophytic lifestyles based on interaction strategies with host plants. *Frontiers in Plant Science* **4**, 120.
- Olanrewaju OS, Glick BR, Babalola OO.** 2017. Mechanisms of action of plant growth promoting bacteria. *World Journal of Microbiology and Biotechnology* **33**, 197.
- Omics.** 2017. Nutrient Film Technique.

**Paradiso R.** 2017. Changes in Leaf Anatomical Traits Enhanced Photosynthetic Activity of Soybean Grown in Hydroponics with Plant Growth-Promoting Microorganisms. *Frontiers in Plant Science* **8**, 674.

**Paradiso R, Buonomo R, Dizon R, Barbieri G, De Pascale S.** 2015. Effect of bacterial root symbiosis and urea as source of nitrogen on performance of soybean plants grown hydroponically for bioregenerative life support systems (BLSSs). *Frontiers in Plant Science* **6**, 888.

**Paradiso R.** 2014. Soilless cultivation of soybean for Bioregenerative Life Support Systems (BLSSs): A literature review and the experience of the MELISSA Project—Food characterization Phase I. *Plant Biology* **16**, 69-78.

**Peer RV, Schippers B.** 1989. Plant growth responses to bacterization with selected *Pseudomonas* spp. Strains and rhizosphere microbial development in hydroponic cultures. *Canadian Journal of Microbiology* **35**, 456-463.

**Phibunwatthanawong T, Riddech N.** 2019. Liquid organic fertilizer production for growing vegetables under hydroponic condition. *International Journal of Recycling of Organic Waste in Agriculture* **8**, 369-380.

**Pignata G, Casale M, Nicola S.** 2017. Water and nutrient supply in horticultural crops grown in soilless culture: Resource efficiency in dynamic and intensive systems. *Advances in research on fertilization Management of Vegetable Crops*. Springer 183-219.

**Pii Y, Graf H, Valentinuzzi F, Cesco S, Mimmo T.** 2018. The effects of plant growth-promoting rhizobacteria (PGPR) on the growth and quality of strawberries. *Acta Horticulturae* **1217**, 231-238.

**Pii Y.** 2015. Microbial interactions in the rhizosphere: Beneficial influences of plant growth-promoting rhizobacteria on nutrient acquisition process-A review. *Biology and Fertility of Soils* **51**, 403-415.

**Prathap M, Ranjitha KBD.** 2015. A Critical Review on Plant Growth Promoting Rhizobacteria. *Journal of Plant Pathology and Microbiology* **6**, 266.

**Saha M.** 2016. Microbial siderophores and their potential applications: A review. *Environmental Science Pollution Research* **23**, 3984-3999.

**Sambo P.** 2019. Hydroponic Solutions for Soilless Production Systems: Issues and Opportunities in a Smart Agriculture Perspective. *Frontiers in Plant Science* **10**, 923-923.

**Santoyo G, Moreno-Hagelsieb G, Orozco-Mosqueda MC, Glick BR.** 2016. Plant growth-promoting bacterial endophytes. *Microbiological Research* **183**, 92-99.

**Selma MV.** 2012. Quality, bioactive constituents and micquality of green and red fresh-cut lettuce (*Lactuca sativa* L.) are influenced by soil agricultural production systems. *Biology and Technology* **63**, 16-24.

**Seungjun L, Jiyoung L.** 2015. Beneficial bacteria and fungi in hydroponic systems: Types and characteristics of hydroponic food production methods. *Scientia Horticulturae* **195**, 206-215.

**Sgherri C, Cecconami S, Pinzino C, Navari-Izzo F, Izzo R.** 2010. Levels of antioxidants and nutraceuticals in basil grown in hydroponics and soil. *Food Chemistry* **123**, 416-422.

**Shinohara M.** 2011. Microbial mineralization of organic nitrogen into nitrate to allow the use of organic fertilizer in hydroponics. *Soil Science and Plant Nutrition* **57**, 190-203.

**Singh M.** 2019. Plant Growth Promoting Rhizobacteria. In *PGPR Amelioration in Sustainable Agriculture*. Elsevier 41-66.

**Song W.** 2004. Tomato Fusarium wilt and its chemical control strategies in a hydroponic system. *Crop Protection* **22**, 243-247.

**Tahmasbi F, Lakzian A, Khavazi K, Pardin A.** 2014. Isolation, Identification and Evaluation of Siderophore Production in Pseudomonas Bacteria and its effect on Hydroponically Grown Con. Journal of Molecular and Cellular Research **27**, 75-87.

**Tank N, Rajendran N, Patel B, Saraf M.** 2012. Evaluation and biochemical characterization of a distinctive pyoverdine from a Pseudomonas isolated from chickpea rhizosphere. Brazilian Journal of Microbiology **43**, 639-648.

**Teotia P, Kumar V, Kumar M, Shrivastava N, Varma A.** 2016. Rhizosphere microbes: Potassium solubilization and crop productivity-present and future aspects. Potassium Solubilizing microorganisms for sustainable agriculture. Springer 315-32.

**Vacheron J.** 2015. Plant growth promoting rhizobacteria and root system functioning. Journal of Soil Science and Plant Nutrition **10**, 293-319.

**Valentinuzzi F.** 2015. Phosphorus and iron deficiencies induce a metabolic reprogramming and affect the exudation traits of the woody plant Fragaria ananassa. Journal of Experimental Botany **66**, 6483-6495.

**Vejan P, Abdullah R, Khadiran T, Ismail S, Boyce AN.** 2016. Role of plant growth promoting rhizobacteria in agricultural sustainability. Molecules **21**, 5-17.

**Wilcox GE.** 1982. The future of hydroponics as a Research Method and Plant Production Method. Journal of Plant Nutrition **5**, 1031-1038.

**Zafar M.** 2012. Effect of Plant Growth-Promoting Rhizobacteria on Growth, Nodulation and Nutrient Accumulation of Lentil Under Controlled Conditions. Pedosphere **22**, 848-859.